

Jet geometrical extension of the KCC-invariants

Vladimir Balan and Mircea Neagu

June 2009; Last revised December 2009 (an added bibliographical item)

Abstract

In this paper we construct the jet geometrical extensions of the KCC-invariants, which characterize a given second-order system of differential equations on the 1-jet space $J^1(\mathbb{R}, M)$. A generalized theorem of characterization of our jet geometrical KCC-invariants is also presented.

Mathematics Subject Classification (2000): 58B20, 37C10, 53C43.

Key words and phrases: 1-jet spaces, temporal and spatial semisprays, nonlinear connections, SODEs, jet h -KCC-invariants.

1 Geometrical objects on 1-jet spaces

We remind first several differential geometrical properties of the 1-jet spaces. The 1-jet bundle

$$\xi = (J^1(\mathbb{R}, M), \pi_1, \mathbb{R} \times M)$$

is a vector bundle over the product manifold $\mathbb{R} \times M$, having the fibre of type $\mathbb{R}^n$, where n is the dimension of the *spatial* manifold M . If the spatial manifold M has the local coordinates $(x^i)_{i=\overline{1,n}}$, then we shall denote the local coordinates of the 1-jet total space $J^1(\mathbb{R}, M)$ by (t, x^i, x_1^i) ; these transform by the rules [12]

$$\begin{cases} \tilde{t} = \tilde{t}(t) \\ \tilde{x}^i = \tilde{x}^i(x^j) \\ \tilde{x}_1^i = \frac{\partial \tilde{x}^i}{\partial x^j} \frac{dt}{dt} \cdot x_1^j. \end{cases} \quad (1.1)$$

In the geometrical study of the 1-jet bundle, a central role is played by the *distinguished tensors* (d -tensors).

Definition 1.1. A geometrical object $D = \left(D_{1k(1)(l)\dots}^{1i(j)(1)\dots} \right)$ on the 1-jet vector bundle, whose local components transform by the rules

$$D_{1k(1)(u)\dots}^{1i(j)(1)\dots} = \tilde{D}_{1r(1)(s)\dots}^{1p(m)(1)\dots} \frac{dt}{dt} \frac{\partial x^i}{\partial \tilde{x}^p} \left(\frac{\partial x^j}{\partial \tilde{x}^m} \frac{d\tilde{t}}{dt} \right) \frac{d\tilde{t}}{dt} \frac{\partial \tilde{x}^r}{\partial x^k} \left(\frac{\partial \tilde{x}^s}{\partial x^u} \frac{dt}{d\tilde{t}} \right) \dots, \quad (1.2)$$

is called a d -tensor field.

Remark 1.2. The use of parentheses for certain indices of the local components

$$D_{1k(1)(l)\dots}^{1i(j)(1)\dots}$$

of the distinguished tensor field D on the 1-jet space is motivated by the fact that the pair of indices " $\begin{smallmatrix} (j) \\ (1) \end{smallmatrix}$ " or " $\begin{smallmatrix} (1) \\ (l) \end{smallmatrix}$ " behaves like a single index.

Example 1.3. The geometrical object

$$\mathbf{C} = \mathbf{C}_{(1)}^{(i)} \frac{\partial}{\partial x_1^i},$$

where $\mathbf{C}_{(1)}^{(i)} = x_1^i$, represents a d -tensor field on the 1-jet space; this is called the *canonical Liouville d -tensor field* of the 1-jet bundle and is a global geometrical object.

Example 1.4. Let $h = (h_{11}(t))$ be a Riemannian metric on the relativistic time axis $\mathbb{R}$. The geometrical object

$$\mathbf{J}_h = J_{(1)1j}^{(i)} \frac{\partial}{\partial x_1^i} \otimes dt \otimes dx^j,$$

where $J_{(1)1j}^{(i)} = h_{11} \delta_j^i$ is a d -tensor field on $J^1(\mathbb{R}, M)$, which is called the *normalization d -tensor field* of the 1-jet space and is a global geometrical object.

In the Riemann-Lagrange differential geometry of the 1-jet spaces developed in [11], [12] important rôles are also played by geometrical objects as the *temporal* or *spatial semisprays*, together with the *jet nonlinear connections*.

Definition 1.5. A set of local functions $H = \left(H_{(1)1}^{(j)} \right)$ on $J^1(\mathbb{R}, M)$, which transform by the rules

$$2\tilde{H}_{(1)1}^{(k)} = 2H_{(1)1}^{(j)} \left(\frac{dt}{d\tilde{t}} \right)^2 \frac{\partial \tilde{x}^k}{\partial x^j} - \frac{dt}{d\tilde{t}} \frac{\partial \tilde{x}_1^k}{\partial t}, \quad (1.3)$$

is called a *temporal semispray* on $J^1(\mathbb{R}, M)$.

Example 1.6. Let us consider a Riemannian metric $h = (h_{11}(t))$ on the temporal manifold $\mathbb{R}$ and let

$$H_{11}^1 = \frac{h^{11}}{2} \frac{dh_{11}}{dt},$$

where $h^{11} = 1/h_{11}$, be its Christoffel symbol. Taking into account that we have the transformation rule

$$\tilde{H}_{11}^1 = H_{11}^1 \frac{dt}{d\tilde{t}} + \frac{d\tilde{t}}{dt} \frac{d^2 t}{d\tilde{t}^2}, \quad (1.4)$$

we deduce that the local components

$$\mathring{H}_{(1)1}^{(j)} = -\frac{1}{2} H_{11}^1 x_1^j$$

define a temporal semispray $\mathring{H} = \left(\mathring{H}_{(1)1}^{(j)} \right)$ on $J^1(\mathbb{R}, M)$. This is called the *canonical temporal semispray associated to the temporal metric $h(t)$* .

Definition 1.7. A set of local functions $G = \left(G_{(1)1}^{(j)} \right)$, which transform by the rules

$$2\tilde{G}_{(1)1}^{(k)} = 2G_{(1)1}^{(j)} \left(\frac{dt}{d\tilde{t}} \right)^2 \frac{\partial \tilde{x}^k}{\partial x^j} - \frac{\partial x^m}{\partial \tilde{x}^j} \frac{\partial \tilde{x}_1^k}{\partial x^m} \tilde{x}_1^j, \quad (1.5)$$

is called a *spatial semispray* on $J^1(\mathbb{R}, M)$.

Example 1.8. Let $\varphi = (\varphi_{ij}(x))$ be a Riemannian metric on the spatial manifold M and let us consider

$$\gamma_{jk}^i = \frac{\varphi^{im}}{2} \left(\frac{\partial \varphi_{jm}}{\partial x^k} + \frac{\partial \varphi_{km}}{\partial x^j} - \frac{\partial \varphi_{jk}}{\partial x^m} \right)$$

its Christoffel symbols. Taking into account that we have the transformation rules

$$\tilde{\gamma}_{qr}^p = \gamma_{jk}^i \frac{\partial \tilde{x}^p}{\partial x^i} \frac{\partial x^j}{\partial \tilde{x}^q} \frac{\partial x^k}{\partial \tilde{x}^r} + \frac{\partial \tilde{x}^p}{\partial x^l} \frac{\partial^2 x^l}{\partial \tilde{x}^q \partial \tilde{x}^r}, \quad (1.6)$$

we deduce that the local components

$$\mathring{G}_{(1)1}^{(j)} = \frac{1}{2} \gamma_{kl}^j x_1^k x_1^l$$

define a spatial semispray $\mathring{G} = \left(\mathring{G}_{(1)1}^{(j)} \right)$ on $J^1(\mathbb{R}, M)$. This is called the *canonical spatial semispray associated to the spatial metric $\varphi(x)$* .

Definition 1.9. A set of local functions $\Gamma = \left(M_{(1)1}^{(j)}, N_{(1)i}^{(j)} \right)$ on $J^1(\mathbb{R}, M)$, which transform by the rules

$$\tilde{M}_{(1)1}^{(k)} = M_{(1)1}^{(j)} \left(\frac{dt}{d\tilde{t}} \right)^2 \frac{\partial \tilde{x}^k}{\partial x^j} - \frac{dt}{d\tilde{t}} \frac{\partial \tilde{x}_1^k}{\partial t} \quad (1.7)$$

and

$$\tilde{N}_{(1)l}^{(k)} = N_{(1)i}^{(j)} \frac{dt}{d\tilde{t}} \frac{\partial x^i}{\partial \tilde{x}^l} \frac{\partial \tilde{x}^k}{\partial x^j} - \frac{\partial x^m}{\partial \tilde{x}^l} \frac{\partial \tilde{x}_1^k}{\partial x^m}, \quad (1.8)$$

is called a *nonlinear connection* on the 1-jet space $J^1(\mathbb{R}, M)$.

Example 1.10. Let us consider that $(\mathbb{R}, h_{11}(t))$ and $(M, \varphi_{ij}(x))$ are Riemannian manifolds having the Christoffel symbols $H_{11}^1(t)$ and $\gamma_{jk}^i(x)$. Then, using the transformation rules (1.1), (1.4) and (1.6), we deduce that the set of local functions

$$\mathring{\Gamma} = \left(\mathring{M}_{(1)1}^{(j)}, \mathring{N}_{(1)i}^{(j)} \right),$$

where

$$\mathring{M}_{(1)1}^{(j)} = -H_{11}^1 x_1^j \quad \text{and} \quad \mathring{N}_{(1)i}^{(j)} = \gamma_{im}^j x_1^m,$$

represents a nonlinear connection on the 1-jet space $J^1(\mathbb{R}, M)$. This jet nonlinear connection is called the *canonical nonlinear connection attached to the pair of Riemannian metrics $(h(t), \varphi(x))$* .

In the sequel, let us study the geometrical relations between *temporal* or *spatial semisprays* and *nonlinear connections* on the 1-jet space $J^1(\mathbb{R}, M)$. In this direction, using the local transformation laws (1.3), (1.7) and (1.1), respectively the transformation laws (1.5), (1.8) and (1.1), by direct local computation, we find the following geometrical results:

Theorem 1.11. *a) The temporal semisprays $H = (H_{(1)1}^{(j)})$ and the sets of temporal components of nonlinear connections $\Gamma_{\text{temporal}} = (M_{(1)1}^{(j)})$ are in one-to-one correspondence on the 1-jet space $J^1(\mathbb{R}, M)$, via:*

$$M_{(1)1}^{(j)} = 2H_{(1)1}^{(j)}, \quad H_{(1)1}^{(j)} = \frac{1}{2}M_{(1)1}^{(j)}.$$

b) The spatial semisprays $G = (G_{(1)1}^{(j)})$ and the sets of spatial components of nonlinear connections $\Gamma_{\text{spatial}} = (N_{(1)k}^{(j)})$ are connected on the 1-jet space $J^1(\mathbb{R}, M)$, via the relations:

$$N_{(1)k}^{(j)} = \frac{\partial G_{(1)1}^{(j)}}{\partial x_1^k}, \quad G_{(1)1}^{(j)} = \frac{1}{2}N_{(1)m}^{(j)}x_1^m.$$

2 Jet geometrical KCC-theory

In this Section we generalize on the 1-jet space $J^1(\mathbb{R}, M)$ the basics of the KCC-theory ([1], [4], [6], [13]). In this respect, let us consider on $J^1(\mathbb{R}, M)$ a second-order system of differential equations of local form

$$\frac{d^2x^i}{dt^2} + F_{(1)1}^{(i)}(t, x^k, x_1^k) = 0, \quad i = \overline{1, n}, \quad (2.1)$$

where $x_1^k = dx^k/dt$ and the local components $F_{(1)1}^{(i)}(t, x^k, x_1^k)$ transform under a change of coordinates (1.1) by the rules

$$\tilde{F}_{(1)1}^{(r)} = F_{(1)1}^{(j)} \left(\frac{dt}{d\tilde{t}} \right)^2 \frac{\partial \tilde{x}^r}{\partial x^j} - \frac{dt}{d\tilde{t}} \frac{\partial \tilde{x}_1^r}{\partial t} - \frac{\partial x^m}{\partial \tilde{x}^j} \frac{\partial \tilde{x}_1^r}{\partial x^m} \tilde{x}_1^j. \quad (2.2)$$

Remark 2.1. The second-order system of differential equations (2.1) is invariant under a change of coordinates (1.1).

Using a temporal Riemannian metric $h_{11}(t)$ on $\mathbb{R}$ and taking into account the transformation rules (1.3) and (1.5), we can rewrite the SODEs (2.1) in the following form:

$$\frac{d^2x^i}{dt^2} - H_{11}^1 x_1^i + 2G_{(1)1}^{(i)}(t, x^k, x_1^k) = 0, \quad i = \overline{1, n},$$

where

$$G_{(1)1}^{(i)} = \frac{1}{2}F_{(1)1}^{(i)} + \frac{1}{2}H_{11}^1 x_1^i$$

are the components of a spatial semispray on $J^1(\mathbb{R}, M)$. Moreover, the coefficients of the spatial semispray $G_{(1)1}^{(i)}$ produce the spatial components $N_{(1)j}^{(i)}$ of a nonlinear connection Γ on the 1-jet space $J^1(\mathbb{R}, M)$, by putting

$$N_{(1)j}^{(i)} = \frac{\partial G_{(1)1}^{(i)}}{\partial x_1^j} = \frac{1}{2} \frac{\partial F_{(1)1}^{(i)}}{\partial x_1^j} + \frac{1}{2} H_{11}^1 \delta_j^i.$$

In order to find the basic jet differential geometrical invariants of the system (2.1) (see Kosambi [10], Cartan [8] and Chern [9]) under the jet coordinate transformations (1.1), we define the *h -KCC-covariant derivative of a d -tensor of kind $T_{(1)}^{(i)}(t, x^k, x_1^k)$* on the 1-jet space $J^1(\mathbb{R}, M)$ via

$$\begin{aligned} \frac{{}^h D T_{(1)}^{(i)}}{dt} &= \frac{dT_{(1)}^{(i)}}{dt} + N_{(1)r}^{(i)} T_{(1)}^{(r)} - H_{11}^1 T_{(1)}^{(i)} = \\ &= \frac{dT_{(1)}^{(i)}}{dt} + \frac{1}{2} \frac{\partial F_{(1)1}^{(i)}}{\partial x_1^r} T_{(1)}^{(r)} - \frac{1}{2} H_{11}^1 T_{(1)}^{(i)}, \end{aligned}$$

where the Einstein summation convention is used throughout.

Remark 2.2. The *h -KCC-covariant derivative* components $\frac{{}^h D T_{(1)}^{(i)}}{dt}$ transform under a change of coordinates (1.1) as a d -tensor of type $\mathcal{T}_{(1)1}^{(i)}$.

In such a geometrical context, if we use the notation $x_1^i = dx^i/dt$, then the system (2.1) can be rewritten in the following distinguished tensorial form:

$$\begin{aligned} \frac{{}^h D x_1^i}{dt} &= -F_{(1)1}^{(i)}(t, x^k, x_1^k) + N_{(1)r}^{(i)} x_1^r - H_{11}^1 x_1^i = \\ &= -F_{(1)1}^{(i)} + \frac{1}{2} \frac{\partial F_{(1)1}^{(i)}}{\partial x_1^r} x_1^r - \frac{1}{2} H_{11}^1 x_1^i, \end{aligned}$$

Definition 2.3. The distinguished tensor

$$\varepsilon_{(1)1}^{(i)} = -F_{(1)1}^{(i)} + \frac{1}{2} \frac{\partial F_{(1)1}^{(i)}}{\partial x_1^r} x_1^r - \frac{1}{2} H_{11}^1 x_1^i$$

is called the *first h -KCC-invariant* on the 1-jet space $J^1(\mathbb{R}, M)$ of the SODEs (2.1), which is interpreted as an *external force* [1], [6].

Example 2.4. It can be easily seen that for the particular first order jet rheonomic dynamical system

$$\frac{dx^i}{dt} = X_{(1)}^{(i)}(t, x^k) \Rightarrow \frac{d^2 x^i}{dt^2} = \frac{\partial X_{(1)}^{(i)}}{\partial t} + \frac{\partial X_{(1)}^{(i)}}{\partial x^m} x_1^m, \quad (2.3)$$

where $X_{(1)}^{(i)}(t, x)$ is a given d -tensor on $J^1(\mathbb{R}, M)$, the first h -KCC-invariant has the form

$$\varepsilon_{(1)1}^{(i)} = \frac{\partial X_{(1)}^{(i)}}{\partial t} + \frac{1}{2} \frac{\partial X_{(1)}^{(i)}}{\partial x^r} x_1^r - \frac{1}{2} H_{11}^1 x_1^i.$$

In the sequel, let us vary the trajectories $x^i(t)$ of the system (2.1) by the nearby trajectories $(\bar{x}^i(t, s))_{s \in (-\varepsilon, \varepsilon)}$, where $\bar{x}^i(t, 0) = x^i(t)$. Then, considering the *variation d -tensor field*

$$\xi^i(t) = \left. \frac{\partial \bar{x}^i}{\partial s} \right|_{s=0},$$

we get the *variational equations*

$$\frac{d^2 \xi^i}{dt^2} + \frac{\partial F_{(1)1}^{(i)}}{\partial x^j} \xi^j + \frac{\partial F_{(1)1}^{(i)}}{\partial x_1^r} \frac{d\xi^r}{dt} = 0. \quad (2.4)$$

In order to find other jet geometrical invariants for the system (2.1), we also introduce the *h -KCC-covariant derivative of a d -tensor of kind $\xi^i(t)$* on the 1-jet space $J^1(\mathbb{R}, M)$ via

$$\frac{{}^h D \xi^i}{dt} = \frac{d\xi^i}{dt} + N_{(1)m}^{(i)} \xi^m = \frac{d\xi^i}{dt} + \frac{1}{2} \frac{\partial F_{(1)1}^{(i)}}{\partial x_1^m} \xi^m + \frac{1}{2} H_{11}^1 \xi^i.$$

Remark 2.5. The *h -KCC-covariant derivative components $\frac{{}^h D \xi^i}{dt}$* transform under a change of coordinates (1.1) as a d -tensor $T_{(1)}^{(i)}$.

In this geometrical context, the variational equations (2.4) can be rewritten in the following distinguished tensorial form:

$$\frac{{}^h D}{dt} \left[\frac{{}^h D \xi^i}{dt} \right] = P_{m11}^i \xi^m,$$

where

$$\begin{aligned} {}^h P_{j11}^i &= -\frac{\partial F_{(1)1}^{(i)}}{\partial x^j} + \frac{1}{2} \frac{\partial^2 F_{(1)1}^{(i)}}{\partial t \partial x_1^j} + \frac{1}{2} \frac{\partial^2 F_{(1)1}^{(i)}}{\partial x^r \partial x_1^j} x_1^r - \frac{1}{2} \frac{\partial^2 F_{(1)1}^{(i)}}{\partial x_1^r \partial x_1^j} F_{(1)1}^{(r)} + \\ &+ \frac{1}{4} \frac{\partial F_{(1)1}^{(i)}}{\partial x_1^r} \frac{\partial F_{(1)1}^{(r)}}{\partial x_1^j} + \frac{1}{2} \frac{dH_{11}^1}{dt} \delta_j^i - \frac{1}{4} H_{11}^1 H_{11}^1 \delta_j^i. \end{aligned}$$

Definition 2.6. The d -tensor ${}^h P_{j11}^i$ is called the *second h -KCC-invariant* on the 1-jet space $J^1(\mathbb{R}, M)$ of the system (2.1), or the *jet h -deviation curvature d -tensor*.

Example 2.7. If we consider the second-order system of differential equations of the *harmonic curves associated to the pair of Riemannian metrics* $(h_{11}(t), \varphi_{ij}(x))$, system which is given by (see the Examples 1.6 and 1.8)

$$\frac{d^2 x^i}{dt^2} - H_{11}^1(t) \frac{dx^i}{dt} + \gamma_{jk}^i(x) \frac{dx^j}{dt} \frac{dx^k}{dt} = 0,$$

where $H_{11}^1(t)$ and $\gamma_{jk}^i(x)$ are the Christoffel symbols of the Riemannian metrics $h_{11}(t)$ and $\varphi_{ij}(x)$, then the second h -KCC-invariant has the form

$$P_{j11}^i = -R_{pqj}^i x_1^p x_1^q,$$

where

$$R_{pqj}^i = \frac{\partial \gamma_{pq}^i}{\partial x^j} - \frac{\partial \gamma_{pj}^i}{\partial x^q} + \gamma_{pq}^r \gamma_{rj}^i - \gamma_{pj}^r \gamma_{rq}^i$$

are the components of the curvature of the spatial Riemannian metric $\varphi_{ij}(x)$. Consequently, the variational equations (2.4) become the following *jet Jacobi field equations*:

$$\frac{h}{dt} \left[\frac{h}{dt} \xi^i \right] + R_{pqm}^i x_1^p x_1^q \xi^m = 0,$$

where

$$\frac{h}{dt} \xi^i = \frac{d\xi^i}{dt} + \gamma_{jm}^i x_1^j \xi^m.$$

Example 2.8. For the particular first order jet rheonomic dynamical system (2.3) the jet h -deviation curvature d -tensor is given by

$$P_{j11}^i = \frac{1}{2} \frac{\partial^2 X_{(1)}^{(i)}}{\partial t \partial x^j} + \frac{1}{2} \frac{\partial^2 X_{(1)}^{(i)}}{\partial x^j \partial x^r} x_1^r + \frac{1}{4} \frac{\partial X_{(1)}^{(i)}}{\partial x^r} \frac{\partial X_{(1)}^{(r)}}{\partial x^j} + \frac{1}{2} \frac{dH_{11}^1}{dt} \delta_j^i - \frac{1}{4} H_{11}^1 H_{11}^1 \delta_j^i.$$

Definition 2.9. The distinguished tensors

$$R_{jk1}^i = \frac{1}{3} \left[\frac{\partial P_{j11}^i}{\partial x_1^k} - \frac{\partial P_{k11}^i}{\partial x_1^j} \right], \quad B_{jkm}^i = \frac{\partial R_{jk1}^i}{\partial x_1^m}$$

and

$$D_{jkm}^{i1} = \frac{\partial^3 F_{(1)1}^{(i)}}{\partial x_1^j \partial x_1^k \partial x_1^m}$$

are called the *third*, *fourth* and *fifth* h -KCC-invariant on the 1-jet vector bundle $J^1(\mathbb{R}, M)$ of the system (2.1).

Remark 2.10. Taking into account the transformation rules (2.2) of the components $F_{(1)1}^{(i)}$, we immediately deduce that the components D_{jkm}^{i1} behave like a d -tensor.

Example 2.11. For the first order jet rheonomic dynamical system (2.3) the third, fourth and fifth h -KCC-invariants are zero.

Theorem 2.12 (of characterization of the jet h -KCC-invariants). *All the five h -KCC-invariants of the system (2.1) cancel on $J^1(\mathbb{R}, M)$ if and only if there exists a flat symmetric linear connection $\Gamma_{jk}^i(x)$ on M such that*

$$F_{(1)1}^{(i)} = \Gamma_{pq}^i(x)x_1^p x_1^q - H_{11}^1(t)x_1^i. \quad (2.5)$$

Proof. " $\Leftarrow$ " By a direct calculation, we obtain

$$\varepsilon_{(1)1}^{h(i)} = 0, \quad \overset{h}{P}_{j11}^i = -\mathfrak{R}_{pqj}^i x_1^p x_1^q = 0 \text{ and } D_{jkl}^{i1} = 0,$$

where $\mathfrak{R}_{pqj}^i = 0$ are the components of the curvature of the flat symmetric linear connection $\Gamma_{jk}^i(x)$ on M .

" $\Rightarrow$ " By integration, the relation

$$D_{jkl}^{i1} = \frac{\partial^3 F_{(1)1}^{(i)}}{\partial x_1^j \partial x_1^k \partial x_1^l} = 0$$

subsequently leads to

$$\begin{aligned} \frac{\partial^2 F_{(1)1}^{(i)}}{\partial x_1^j \partial x_1^k} &= 2\Gamma_{jk}^i(t, x) \Rightarrow \frac{\partial F_{(1)1}^{(i)}}{\partial x_1^j} = 2\Gamma_{jp}^i x_1^p + \mathcal{U}_{(1)j}^{(i)}(t, x) \Rightarrow \\ &\Rightarrow F_{(1)1}^{(i)} = \Gamma_{pq}^i x_1^p x_1^q + \mathcal{U}_{(1)p}^{(i)} x_1^p + \mathcal{V}_{(1)1}^{(i)}(t, x), \end{aligned}$$

where the local functions $\Gamma_{jk}^i(t, x)$ are symmetrical in the indices j and k .

The equality $\varepsilon_{(1)1}^{h(i)} = 0$ on $J^1(\mathbb{R}, M)$ leads us to

$$\mathcal{V}_{(1)1}^{(i)} = 0$$

and to

$$\mathcal{U}_{(1)j}^{(i)} = -H_{11}^1 \delta_j^i.$$

Consequently, we have

$$\frac{\partial F_{(1)1}^{(i)}}{\partial x_1^j} = 2\Gamma_{jp}^i x_1^p - H_{11}^1 \delta_j^i$$

and

$$F_{(1)1}^{(i)} = \Gamma_{pq}^i x_1^p x_1^q - H_{11}^1 x_1^i.$$

The condition $\overset{h}{P}_{j11}^i = 0$ on $J^1(\mathbb{R}, M)$ implies the equalities $\Gamma_{jk}^i = \Gamma_{jk}^i(x)$ and

$$\mathfrak{R}_{pqj}^i + \mathfrak{R}_{qpj}^i = 0,$$

where

$$\mathfrak{R}_{pqj}^i = \frac{\partial \Gamma_{pq}^i}{\partial x^j} - \frac{\partial \Gamma_{pj}^i}{\partial x^q} + \Gamma_{pq}^r \Gamma_{rj}^i - \Gamma_{pj}^r \Gamma_{rq}^i.$$

It is important to note that, taking into account the transformation laws (2.2), (1.3) and (1.1), we deduce that the local coefficients $\Gamma_{jk}^i(x)$ behave like a symmetric linear connection on M . Consequently, $\mathfrak{R}_{pqj}^i$ represent the curvature of this symmetric linear connection.

On the other hand, the equality $\overset{h}{R}_{jk1}^i = 0$ leads us to $\mathfrak{R}_{qjk}^i = 0$, which infers that the symmetric linear connection $\Gamma_{jk}^i(x)$ on M is flat. $\square$

Acknowledgements. The present research was supported by the Romanian Academy Grant 4/2009.

References

- [1] P.L. Antonelli, *Equivalence Problem for Systems of Second Order Ordinary Differential Equations*, Encyclopedia of Mathematics, Kluwer Academic, Dordrecht, 2000.
- [2] P.L. Antonelli, L. Bevilacqua, S.F. Rutz, *Theories and models in symbiogenesis*, Nonlinear Analysis: Real World Applications **4** (2003), 743-753.
- [3] P.L. Antonelli, I. Bucătaru, *New results about the geometric invariants in KCC-theory*, An. Șt. Univ. "A.I.Cuza" Iași. Mat. N.S. **47** (2001), 405-420.
- [4] P.L. Antonelli, R.S. Ingarden, M. Matsumoto, *The Theory of Sprays and Finsler Spaces with Applications in Physics and Biology*, Fundamental Theories of Physics, vol. **58**, Kluwer Academic Publishers, Dordrecht, 1993.
- [5] Gh. Atanasiu, V. Balan, N. Brînzei, M. Rahula, *The Differential-Geometric Structures: Tangent Bundles, Connections in Bundles and The Exponential Law in Jet-Spaces* (in Russian), LKI, Moscow, 2010.
- [6] V. Balan, I.R. Nicola, *Linear and structural stability of a cell division process model*, Hindawi Publishing Corporation, International Journal of Mathematics and Mathematical Sciences, vol. 2006, 1-15.
- [7] I. Bucătaru, R. Miron, *Finsler-Lagrange Geometry. Applications to Dynamical Systems*, Romanian Academy Eds., 2007.
- [8] E. Cartan, *Observations sur le mémoire précédent*, Mathematische Zeitschrift **37** (1933), no. **1**, 619-622.
- [9] S.S. Chern, *Sur la géométrie d'un système d'équations différentielles du second ordre*, Bulletin des Sciences Mathématiques **63** (1939), 206-212.
- [10] D.D. Kosambi, *Parallelism and path-spaces*, Mathematische Zeitschrift **37** (1933), no. **1**, 608-618.

- [11] M. Neagu, *Riemann-Lagrange Geometry on 1-Jet Spaces*, Matrix Rom, Bucharest, 2005.
- [12] M. Neagu, C. Udriște, A. Oană, *Multi-time dependent sprays and h -traceless maps on $J^1(T, M)$* , Balkan Journal of Geometry and Its Applications, vol. **10** (2005), no. **2**, 76-92.
- [13] V.S. Sabău, *Systems biology and deviation curvature tensor*, Nonlinear Analysis. Real World Applications **6** (2005), no. **3**, 563-587.

Authors' addresses:

Vladimir Balan

University Politehnica of Bucharest, Faculty of Applied Sciences,

Department of Mathematics-Informatics I,

Splaiul Independenței 313, RO-060042 Bucharest, Romania.

E-mail: vladimir.balan@upb.ro

Website: <http://www.mathem.pub.ro/dept/vbalan.htm>

Mircea Neagu

University Transilvania of Braşov, Faculty of Mathematics and Informatics,

Department of Algebra, Geometry and Differential Equations,

B-dul Iuliu Maniu, Nr. 50, BV 500091, Braşov, Romania.

E-mail: mircea.neagu@unitbv.ro

Website: <http://www.2collab.com/user/mirceaneagu>